
Intended use:

This product is designed for use under recommendations from your health care professional or sleep therapist. This product is used on adult patients (weight above 30kg) whom CPAP or bi-level therapy has been prescribed. The product is re-usable for single patient in medical institutions, home care and sleep treatment center.



Instructions for use:

- ➤ Inspect the packaging and product appearance before using. Please do not use if there are any worn or damaged parts.
- Take out the mask set and L-type connector from the box and snap the L-type connector into the mask.
- > See headgear assembly diagram, hook headgear point A to the mask frame connecting rings on both sides, and then bond the two B points.
- > Hold the mask over nose and mouth, and then position the headgear.
- Fast release buckle to the mask frame Quick release buckle connection column.
- Adjust the head straps and the hook and loop of the mask comfortably.
- ➤ L-type connector side ISO 4.3mmOD hole for measurement monitoring use.
- > Connect the measuring tube /smoothbore tube to use for needed users.
- After wearing, connect the ventilation tube connector to the ISO 22mm OD for use.
- Quick release buckle on both sides can be easily removed, also be connected easily.

♦ Cleaning:

- ♦ This product is reusable for single patient.
- → Follow the procedures for cleaning/sterilization in medical institutions.
- Wash the headgear with warm water and neutral detergent or soap. After rinse, air-dry the strap and keep away from direct sunlight or the dryer.
- Mask and L-type connector can be disconnected for cleaning.
- Wash with warm water and neutral detergent or soap.
- ♦ After rinse, dry in a cool place. Do not put it in the dryer.
- ♦ The uncovered can be autoclaved 25 times with the 15minute*121°C high temperature steam (pressure) sterilization method.



Warning:

- \triangle Multiple patients reuse will result in cross-infection of disease.
- △ Prohibit the use of if the patient has the following symptoms: vomiting, facial trauma burns, impaired myocardial function, Gastroesophageal Reflux Disease, abnormal cough reflex, unconsciousness.
- \triangle To be worn by professional medical staff guidance.
- \triangle Do not cover or block anti-asphyxiation holes and breathing valve.
- \triangle Do not connect worn breathing apparatus has not been started before operation.
- \triangle If you feel unwell, STOP using it immediately and seek medical advice immediately.
- \triangle Do not use alcohol \cdot bleach or solvents containing chlorine and other components to clean the mask set.
- △ Do not put the mask set and headgear in the place of the sun exposure or high temperature preservation.
- △ Do not dismantling the silicone mask and, this action will mask a small loss of functionality, and reduce product life.
- Occur in a leak, slight tightening of the headgear, no improvement in the condition of leakage and then gently tighten the headgear, do not force too much cause discomfort.
- \triangle Dispose of it according to the local medical waste disposal regulations.

Specifications and materials :

- Mask size : S / M / L
- L-type connector: 22mmOD meet ISO 5356-1.
- Headgear : General.
- Measuring tube: 6mmOD, 3.8mmID, 200cm.
- Smoothbore tube: 180cm, 22mmID meet ISO 5356-1.

Mask size	LARGE			MEDIUM			SMALL	
Dead space	172 mL			164 mL			152 mL	
Therapy pressure	4 to 20 cm H ₂ O							
Sound power	Average: 31.2 db(A)							
Sound pressure	Average: 22.5 db(A)							
Connector	22mmOD meet ISO 5356-1							
Measuring hole	4.3mmOD meet EN 20594-1							
Resistance	50 L/min : 0.22 cmH₂O, 100 L/min : 0.46 cmH₂O							
Exhaust Flow	Pressure (cmH ₂ O)		5	10	15	20	25	30
Rate at Given	Flow (L/min)		23	35	47	56	68	80
Therapy Pressure	Flow(L/min)	80 ————————————————————————————————————	Pressure-flow curve					
		4	5	10 Pressure	15 (cmH2O)	20	25	30





(€ 2460





Obelis s.a. Boulevard Général Wahis 53 1030 Brussels, BELGIUM Tel: +(32) 2.732.59.54 Fax: +(32)2.732.60.03 E-Mail: mail@obelis.net

